

Performing Physician _____

Arrival Time _____

PM Bowel Preparation with SUPREP

- **You must be accompanied by a friend or relative to drive and assist you home. We WILL NOT discharge you** to a cab, bus, uber or other transportation service **without** a responsible party with you. You may not drive until the day after your procedure.

Special Instructions Regarding Medications

- If you are on an **anticoagulant/blood thinner** our nurse will contact you with instructions after contacting your prescribing physician. If you have not heard from the nurse 1 week prior to your procedure please contact our office.
- **Do Not** take **Iron supplements** for **one week prior** to the procedure date. Multivitamins are OK.
- **Diabetic patients:** _____
- **Do Not** take **Water Pill/Diuretic** (Lasix (furosemide), HCTZ (Hydrochlorothiazide), Dyazide, Diovan HCT, Spironolactone), **on the day of the procedure**
- **Heart/Blood Pressure/Thyroid/Seizure/Anxiety Medications** can be taken **on the day of the procedure** with a sip of water.
- **Stop Effexor one day before** the procedure (if applicable)
- **Asthmatic patients** should bring their **INHALERS** with them on the day of the procedure
- **Do Not use Cannabis 12 hours prior to procedure**
- **All Facial Piercings MUST be removed prior to procedure**

THE DAY BEFORE YOUR PROCEDURE

IMPORTANT NOTE: follow the instructions below, NOT the instructions on the prep's packaging

- **You MAY Eat a light breakfast until 11:00 am**
(ex: scrambled eggs, plain white toast, orange juice. **NO** whole grains, cereals, oatmeal, etc.)
- **YOU MAY NOT HAVE ANY SOLID FOOD AFATER 11:00 am ONLY CLEAR LIQUIDS**
- Take all of your normal prescription medications, **unless advised otherwise**.
- **At 6 pm** - yes no Take 2 Dulcolax (Bisacodyl) Laxative tablets with water **2** hours before first dose of prep
- **At 8 pm** – Pour (1) 6 oz bottle of **Suprep** liquid into the mixing container
Add cool drinking water to the 16 oz line on the container and mix.
- Begin drinking the **Suprep** solution. Drink all the liquid in the container
- You **MUST** drink two (2) more 16 oz containers of clear liquid over the next 2 hours

DAY OF YOUR PROCEDURE

- **At 6 am** -
Pour (1) 6 oz bottle of Suprep liquid into the mixing container
Add cool drinking water to the 16 oz line on the container and mix.
- Drink all the liquid in the container
- You **MUST** drink two (2) more 16 oz containers of clear liquids over the next 2 hours.

All prep and clear liquids MUST be completed by 8:30am. No additional liquids, hard candy, chewing gum after 8:30am.

Prep instructions, patient forms, and FAQ's can also be found on our website:
Westchesterendocenter.com

General or reschedule questions call 610-431-3122